

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 757 099 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
05.02.1997 Bulletin 1997/06

(51) Int Cl.⁶: **C12N 15/12, C07K 14/705,
C12N 15/85, C12N 5/10,
A61K 38/17**

(21) Application number: **96305298.0**

(22) Date of filing: **19.07.1996**

(84) Designated Contracting States:
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE**

- **Ledbetter, Jeffrey A.**
Seattle, WA 98117 (US)
- **Peach, Robert**
Edmonds, WA 98026 (US)

(30) Priority: **21.07.1995 US 505058**

(71) Applicant: **Bristol-Myers Squibb Company**
New York, N.Y. 10154 (US)

(74) Representative: **Jones, Alan John et al**
CARMAELS & RANSFORD
43 Bloomsbury Square
London, WC1A 2RA (GB)

(72) Inventors:
• **Linsley, Peter S.**
Seattle, WA 98119 (US)

Remarks:

The applicant has subsequently filed a sequence listing and declared, that it includes no new matter.


(54) CTLA4 mutant molecules and uses thereof

(57) The invention provides CTLA4 mutant molecules as ligands for the B7 antigen. Methods are provided for expressing CTLA4 mutant molecules as soluble,

functional molecules, for preparing CTLA4 mutant fusion proteins, and for using these soluble molecules to regulate T cell interactions and immune responses mediated by such interactions.

FIGURE 7

Wild type (Wt.) and Mutant Binding to CD80 CHO and CD86 CHO
as determined by FACS

**EP 0 757 099 A2**

Description

Throughout this application various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

The present invention relates to expression of CTLA4 mutant molecules and to methods for regulating cellular interactions using the CTLA4 mutant molecules.

BACKGROUND OF THE INVENTION

Central events leading to a specific immune response to antigen involve the intimate association of T lymphocytes with antigen presenting cells (APC). In order for T-cells to successfully mount cell-mediated or antibody-mediated immune responses to an antigenic stimulus, distinct activation signals are required from the APC. An antigen-specific signal occurs when the T-cell antigen receptor binds to antigenic peptides presented by major histocompatibility complex (MHC) molecules on the surface of APC's. However, this event alone is not sufficient to stimulate T-cells to proliferate and, by itself, can lead to clonal inactivation or anergy (Schwartz, R.H. (1990) *Science* **248**:1349-1356). For an effective immune response to occur, a second non-antigen-specific co-stimulatory signal must also be delivered to the T-cell from an APC (Freeman et al. (*J. Immunol.* **143**(8):2714-2722 (1989)).

CTLA4 is a T-cell surface receptor that associates with the B7 counter receptor, namely, B7-1 (CD80) and B7-2 (CD86), expressed on antigen-presenting cells (Hathcock et al. "Comparative Analysis of B7-1 and B7-2 Co-Stimulatory Ligands: Expression and Function" 1994 *Journal of Experimental Medicine*, **180**(2):631-40). This association establishes the molecular basis for an important T-cell co-stimulatory pathway, the primary function of which is to induce T-cell cytokine production and proliferation following exposure to antigen (Linsley et al., *J. Exp. Med.* **173**:721-730 (1991)). Confirmatory data has shown that one embodiment of CTLA4, namely, CTLA4Ig, is a potent inhibitor of *in vitro* immune responses by blocking CD28/B7 interactions, thereby preventing T-cell proliferation and inducing antigen-specific unresponsiveness (Blazar et al. "In Vivo Blockade of CD28/CTLA4: B7/BB1 Interaction With CTLA4-Ig Reduces Lethal Murine Graft-Versus-Host Disease Across the Major Histocompatibility Complex Barrier in Mice" *Blood*, 1994 June 15, **83**(12):3815-25).

The nature of these co-stimulatory signals has been the focus of intense research efforts, the goals of which have been not only to understand the immune system, but also to develop therapeutic agents which could block the delivery of co-stimulatory signals thereby further manipulating immune responses. The CTLA4 mutant molecules of the invention have been developed to meet these goals.

SUMMARY OF THE INVENTION

The present invention discloses CTLA4 molecules which have been mutated in order to influence their ability to bind the B7 ligands. These mutations are localized to the extracellular domain of CTLA4. Specifically, the mutations are generated at the position of the first tyrosine moiety within the MYPPPY motif of the extracellular domain of CTLA4. We show that the mutants (i.e., those having any one of alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, or valine at this position) have the ability to bind B7-1 (CD80). However, some do not bind B7-2 (CD86).

Additionally provided are methods for making and using the CTLA4 mutant molecules of the invention.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a graph showing the sequencing alignment of CD28 and CTLA4 family members. Sequences of human (H), mouse (M), rat (R), and chicken (Ch) CD28 are aligned with human and mouse CTLA4. Residues are numbered from the mature protein N-terminus with the signal peptides and transmembrane domains underlined and the CDR-analogous regions noted. Dark shaded areas highlight complete conservation of residues while light shaded areas highlight conservative amino acid substitutions in all family members (SEQ ID NOS 1-6).

Figure 2 is a line graph showing CTLA4Ig and CD28Ig mutants bind B7-1.

Figure 3 is a schematic map of CTLA4/CD28Ig hybrid fusion proteins. Open areas represent CD28 sequence; filled areas represent CTLA4 sequence; cross-hatched areas represent beginning of IgG Fc (also refer to Table I).

Figure 4 is a line graph showing that CTLA4/CD28Ig hybrid fusion proteins bind with high avidity to B7-1 CHO cells.

Figure 5 is a line graph showing that CTLA4/CD28Ig hybrid fusion proteins bind with high avidity to B7-1 CHO cells.

Figure 6 is a molecular model of monomeric CTLA4Ig v-like extracellular domain.

Figure 7 is a bar graph comparison of the ability of CTLA4Ig wild type (w.t.) and mutant molecules to bind CHO

cells expressing either CD80 or CD86. The wild type molecule is designated by MYPPPY, the sequence of the naturally occurring motif. The individual mutants are denoted in the same manner. The letter symbol of the amino acid substituted at the second position in this hexapeptide indicates the specific mutation.

Figure 8 is a comparison of wild type (w.t.) and mutant binding ability using different concentrations of these CTLA4Ig fusion proteins. As disclosed in Figure 7, the mutants are indicated by their respective motif sequences.

Figure 9 is an antibody competition study with the LCL 816 cell line. This figure compares CTLA4Ig wild type and mutant protein binding to CD80 or CD86 in the presence of different concentrations of anti-CD80 or anti-CD86 antibody. As disclosed in Figure 7, the specific proteins are indicated by their respective motif sequences.

DETAILED DESCRIPTION OF THE INVENTION

DEFINITION

As used in this application, the following words or phrases have the meanings specified.

As used herein "blocking B7 interaction" means to interfere with the binding of the B7 antigens to their ligands such as CD28 and/or CTLA4 thereby obstructing T cell and B cell interaction.

As used herein a "B7-binding molecule" means any molecule which will bind any one or both of the B7 antigens.

As used herein a "molecule reactive with the CD80 antigen" means any molecule which will recognize and bind CD80.

As used herein a "molecule reactive with the CD86 antigen" means any molecule which will recognize and bind CD86.

As used herein a "non-CTLA4 molecule" means any molecule which can be attached or joined to the extracellular domain of CTLA4 and does not interfere with CTLA4 binding to its target. These molecules include a polypeptide tag, an immunoglobulin (Ig) tail, a biologically or chemically active protein such as the papillomavirus E7 gene product, melanoma-associated antigen p97, and HIV env protein or a sequence of amino acids which renders soluble and active the extracellular portion of CTLA4 or mutagenized forms thereof.

In order that the invention herein described may be more fully understood, the following description is set forth.

COMPOSITIONS OF THE INVENTION

The invention provides CTLA4 mutant molecules reactive with the CD80 antigen, wherein in the extracellular domain of CTLA4 the first tyrosine in the amino acid motif MYPPPY is replaced by an amino acid other than tyrosine. CTLA4 mutant molecules may be embodied in many forms. The only limitation being that it retain its ability to bind CD80.

In one embodiment of the invention, the CTLA4 mutant molecule is a functional, soluble CTLA4 mutant molecule. The extracellular domain of CTLA4 is an example of a soluble CTLA4 mutant molecule. In one example, the soluble CTLA4 mutant molecule binds the CD80 antigen but does not bind the CD86 antigen.

In a further embodiment of the invention, the soluble CTLA4 mutant molecule has the extracellular domain of mutagenized CTLA4 joined to a non-CTLA4 molecule (also referred to as CTLA4 mutant fusion proteins) such as a polypeptide tag. Other embodiments of soluble CTLA4 molecules include those having the extracellular domain of mutagenized CTLA4 fused or joined with a portion of a biologically or chemically active protein such as the papillomavirus E7 gene product, melanoma-associated antigen p97, and HIV env protein.

In accordance with the practice of this invention, CTLA4 mutant and homolog molecules of the invention may have other amino acid substitutions and still retain the functional property of CTLA4 mutant molecules, namely, the molecule having such substitutions will still bind the CD80 antigen. These amino acid substitutions include, but are not necessarily limited to, amino acid substitutions known in the art as "conservative".

For example, it is a well-established principle of protein chemistry that certain amino acid substitutions, entitled "conservative amino acid substitutions," can frequently be made in a protein without altering either the conformation or the function of the protein. Such changes include substituting any of isoleucine (I), valine (V), and leucine (L) for any other of these hydrophobic amino acids; aspartic acid (D) for glutamic acid (E) and vice versa; glutamine (Q) for asparagine (N) and vice versa; and serine (S) for threonine (T) and vice versa. Other substitutions can also be considered conservative, depending on the environment of the particular amino acid and its role in the three-dimensional structure of the protein. For example, glycine (G) and alanine (A) can frequently be interchangeable, as can alanine and valine (V).

Methionine (M), which is relatively hydrophobic, can frequently be interchanged with leucine and isoleucine, and sometimes with valine. Lysine (K) and arginine (R) are frequently interchangeable in locations in which the significant feature of the amino acid residue is its charge and the differing pK's of these two amino acid residues are not significant. Still other changes can be considered "conservative" in particular environments.

METHODS FOR MAKING THE COMPOSITIONS OF THE INVENTION

The techniques for cloning and expressing DNA sequences encoding the amino acid sequences corresponding to the CTLA4 mutant molecules, e.g., synthesis of oligonucleotides, PCR, transforming cells, constructing vectors, expression systems, and the like are well-established in the art, and most practitioners are familiar with the standard resource materials for specific conditions and procedures. However, the following paragraphs are provided for convenience and notation of modifications of standard techniques where necessary, and may serve as a guideline.

Cloning and Expression of Coding Sequences for Receptors and Fusion Proteins

Constructs corresponding to CTLA4 mutant molecules of the present invention were prepared as described by Linsley et al., J. Exp. Med. 173:721-730 (1991), incorporated by reference herein. Alternatively, cDNA clones may be prepared from RNA obtained from cells expressing B7 antigen and CD28 receptor based on knowledge of the published sequences for these proteins (Aruffo and Seed, Proc. Natl. Acad. Sci. 84:8573-8577 (1987), and Freeman et al. "Murine B7-2, an Alternative CTLA4 Counter-Receptor that Co-Stimulates T-cell Proliferation and Interleukin 2 Production" (1993) Journal of Experimental Medicine 178(6) :2185-92) using standard procedures.

CTLA4 mutant molecules consisting of DNA encoding amino acid sequences corresponding to the extracellular domain of mutagenized CTLA4 and the hinge, CH2 and CH3 regions of human IgC γ 1 were constructed by ligation of PCR fragments. The cDNA encoding the amino acid sequences is amplified using the polymerase chain reaction ("PCR") technique (U.S. Patent Nos. 4,683,195 and 4,683,202 to Mullis et al. and Mullis & Faloona, Methods Enzymol. 154:335-350 (1987)). CTLA4Ig mutant fusion polypeptides were obtained having DNA encoding amino acid sequences from about position 1 to about position 125 of the extracellular domain of CTLA4 and corresponding to the hinge, CH2 and CH3 regions of IgC γ 1.

PCR cDNA made from the total cellular RNA of several human leukemia cell lines was screened, using as primers, oligonucleotides from the published sequence of the CTLA4 gene (Dariavach et al., supra). Of the cDNA tested, H38 cells (an HTLV II-associated leukemia line) provided the best yield of PCR products having the expected size. Since a signal peptide for CTLA4 was not identified in the CTLA4 gene, the N terminus of the predicted sequence of CTLA4 was fused to the signal peptide of oncostatin M (Malik et al., Molec. and Cell. Biol. 9:2847 (1989)) in two steps using oligonucleotides. The product of the PCR reaction was ligated with cDNA encoding the amino acid sequences corresponding to the hinge, CH2 and CH3 regions of IgC γ 1 into a expression vector, such as CDM8 or π LN.

To obtain DNA encoding full length human CTLA4, a cDNA encoding the transmembrane and cytoplasmic domains of CTLA4 was obtained by PCR from H38 cells and joined with a fragment from CTLA4Ig, obtained as described above, encoding the oncostatin M signal peptide fused to the N terminus of CTLA4, using oligonucleotide primers as described in the Examples, infra. PCR fragments were ligated into the plasmid CDM8, resulting in an expression plasmid encoding the full length CTLA4 gene, and designated OMCTLA4.

To generate mutations in specific regions of the CTLA4 sequence, site-directed mutagenesis was performed on the π LN vector which contained the soluble chimeric form of CTLA4 (CTLA4Ig) in which the extracellular domain of CTLA4 was genetically fused to the hinge and constant regions of a human IgG heavy chain (Linsley et al, J. Exp. Med. 173:721-730 (1991)). The CTLA4Ig site-directed mutants were prepared by encoding the desired mutation in overlapping oligonucleotide primers and generating the mutants by PCR (Ho et al., 1989, supra.) using the π LN CTLA4Ig plasmid construct as a template.

In order to generate a comprehensive series of mutants which were directed at the extracellular domain, the codon of the first tyrosine in the MYPPPY motif was changed in order to generate new codons encoding each of the other nineteen amino acids. As this motif appears to be crucial for the CTLA-B7 ligand interactions, all CTLA4 molecules which contain these mutations will have an altered ability to bind the B7 antigens.

To produce large quantities of cloned DNA, vectors containing mutant CTLA4 DNA of the invention were transformed into suitable host cells, such as the bacterial cell line E. coli strain MC1061/p3 (Invitrogen Corp., San Diego, CA) using standard procedures, and colonies are screened for the appropriate plasmids.

The clones containing mutant CTLA4 DNA obtained as described above are then transfected into suitable host cells for expression. Depending on the host cell used, transfection is performed using standard techniques appropriate to such cells. For example, transfection into mammalian cells is accomplished using DEAE-dextran mediated transfection, CaPO₄ co-precipitation, lipofection, electroporation, or protoplast fusion, and other methods known in the art including: lysozyme fusion or erythrocyte fusion, scraping, direct uptake, osmotic or sucrose shock, direct microinjection, indirect microinjection such as via erythrocyte-mediated techniques, and/or by subjecting host cells to electric currents. The above list of transfection techniques is not considered to be exhaustive, as other procedures for introducing genetic information into cells will no doubt be developed.

Expression in eukaryotic host cell cultures derived from multicellular organisms is preferred (Tissue Cultures, Academic Press, Cruz and Patterson, Eds. (1973)). These systems have the additional advantage of the ability to splice

out introns and thus can be used directly to express genomic fragments. Useful host cell lines include Chinese hamster ovary (CHO), monkey kidney (COS), VERO and HeLa cells. In the present invention, cell lines stably expressing the fusion constructs are preferred.

Expression vectors for such cells ordinarily include promoters and control sequences compatible with mammalian cells such as, for example, CMV promoter (CDMB vector) and avian sarcoma virus (ASV) (μ LN vector). Other commonly used early and late promoters include those from Simian Virus 40 (SV 40) (Fiers, et al., Nature 273:113 (1973)), or other viral promoters such as those derived from polyoma, Adenovirus 2, and bovine papilloma virus. The controllable promoter, hMTII (Karin, et al., Nature 299:797-802 (1982)) may also be used. General aspects of mammalian cell host system transformations have been described by Axel (U.S. Patent No. 4,399,216 issued Aug. 16, 1983). It now appears, that "enhancer" regions are important in optimizing expression; these are, generally, sequences found upstream or downstream of the promoter region in non-coding DNA regions. Origins of replication may be obtained, if needed, from viral sources. However, integration into the chromosome is a common mechanism for DNA replication in eukaryotes.

Although preferred host cells for expression of the fusion constructs include eukaryotic cells such as COS or CHO cells, other eukaryotic microbes may be used as hosts. Laboratory strains of Saccharomyces cerevisiae, Baker's yeast, are most used although other strains such as Schizosaccharomyces pombe may be used. Vectors employing, for example, the 2 μ origin of replication of Broach, Meth. Enz. 101:307 (1983), or other yeast compatible origins of replications (for example, Stinchcomb et al., Nature 282:39 (1979)); Tschempe et al., Gene 10:157 (1980); and Clarke et al., Meth. Enz. 101:300 (1983)) may be used. Control sequences for yeast vectors include promoters for the synthesis of glycolytic enzymes (Hess et al., J. Adv. Enzyme Reg. 7:149 (1968); Holland et al., Biochemistry 17:4900 (1978)). Additional promoters known in the art include the CMV promoter provided in the CDMB vector (Toyama and Okayama, FEBS 268:217-221 (1990); the promoter for 3-phosphoglycerate kinase (Hitzeman et al., J. Biol. Chem. 255:2073 (1980)), and those for other glycolytic enzymes. Other promoters, which have the additional advantage of transcription controlled by growth conditions are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, and enzymes responsible for maltose and galactose utilization. It is also believed terminator sequences are desirable at the 3' end of the coding sequences. Such terminators are found in the 3' untranslated region following the coding sequences in yeast-derived genes.

Alternatively, prokaryotic cells may be used as hosts for expression. Prokaryotes most frequently are represented by various strains of E. coli; however, other microbial strains may also be used. Commonly used prokaryotic control sequences which are defined herein to include promoters for transcription initiation, optionally with an operator, along with ribosome binding site sequences, include such commonly used promoters as the beta-lactamase (penicillinase) and lactose (lac) promoter systems (Chang et al., Nature 198: 1056 (1977)), the tryptophan (trp) promoter system (Goeddel et al., Nucleic Acids Res. 8:4057 (1980)) and the lambda derived P_L promoter and N-gene ribosome binding site (Shimatake et al., Nature 292:128 (1981)).

The nucleotide sequences encoding CTLA4 mutant molecules may be expressed in a variety of systems as set forth below. The cDNA may be excised by suitable restriction enzymes and ligated into suitable prokaryotic or eukaryotic expression vectors for such expression. The expression of CTLA4 mutant molecules as fusion proteins permits dimer formation of these proteins.

Expression of the mutant CTLA4 receptor of the invention is accomplished transfecting a cell line such as COS cells, and detecting expression by binding of the CTLA4-transfected cells to a ligand for the CTLA4 receptor, for example by testing for binding of the cells to B7Ig fusion protein. Expression of the mutant CTLA4Ig fusion protein of the invention is accomplished by transfecting a cell line such as COS cells, and detecting expression by testing for binding of the mutant CTLA4Ig fusion protein to cells expressing the appropriate ligand.

Sequences of the resulting constructs are confirmed by DNA sequencing using known procedures, for example as described by Sanger et al., Proc. Natl. Acad. Sci. USA 74:5463 (1977), as further described by Messing et al., Nucleic Acids Res. 9:309 (1981), or by the method of Maxam et al. Methods Enzymol. 65:499 (1980)).

Construction and Expression of soluble CTLA4 mutant molecules

The appropriate DNA fragments of CTLA4 mutant molecules and the protein partners or tags (e.g., biologically or chemically active molecules such as ovalbumin, p97, E7, and env gp120) can be isolated from the cDNA by PCR ((U. S. Patent Nos. 4,683,195 and 4,683,202 to Mullis et al. and Mullis & Faloona, Methods Enzymol. 154:335-350 (1987)).

DNA for the fusion proteins of CTLA4 can be prepared by ligating DNA of CTLA4 with that of the various protein tags. In ELISA and FACS assays, the binding and expression of the soluble CTLA4 mutant molecules was detected using antibodies directed against the tag, e.g., ovalbumin, env gp120, HPV E7, and p97.

The DNA sequence of the ovalbumin gene is known (Schweers et al. J. Biol. Chem. (1990) 265(13):7590-5); the DNA sequence of the E7 papillomavirus oncogene is known (Tindle et al. J. Gen. Vir. (1990) 71:1347-54; the DNA sequence of the melanoma-associated antigen p97 is known (Kahn et al. J. Immunol. (1991) 146(9):3235-41); the DNA sequence of env gp120 is known (Wain-Hobson et al. "Nucleotide sequence of AIDS virus LAV" Cell (1985) 40:9-17; Ratner et

al. "Complete nucleotide sequence of the AIDS virus HTLV3" Nature 313:277-284 (1985)). The identity of the resulting genes for each of the soluble molecules (i.e., fusion proteins) can be confirmed by DNA sequencing.

The cDNA of the fusion proteins can be expressed either in mammalian (cos, DEAE dextran transfection) or insect (baculovirus transfection) cell lines (Jones et al. Nature (1986) 523:346). The supernatants of the transfected cell lines were harvested, assayed and the fusion proteins then purified by affinity chromatography.

Recovery of Protein Products

DNA encoding the amino acid sequence corresponding to the extracellular domains of CTLA4, including the codons for a signal sequence such as that of oncostatin M in cells capable of appropriate processing is fused with DNA encoding the amino acid sequence corresponding to the Fc domain of a naturally dimeric protein. Purification of these fusion protein products after secretion from the cells is thus facilitated using antibodies reactive with the anti-immunoglobulin portion of the fusion proteins. When secreted into the medium, the fusion protein product is recovered using standard protein purification techniques, for example by application to protein A columns.

USES OF THE COMPOSITIONS OF THE INVENTION

CTLA4 proteins which have specific amino acid substitutions at the position of the first tyrosine in the MYPPPY motif have unique binding abilities. Mutants which are generated by replacing this tyrosine with for example either phenylalanine or tryptophan have the capability of discriminating between the B7 antigens. Specifically, these mutant molecules will bind the B7-1 (CD80) antigen but not the B7-2 (CD86) antigen. Therefore these mutant molecules may be used to distinguish between these two antigens. In addition, these mutants may be used to differentially affect the specific biological processes which are mediated by the different B7 antigens.

CTLA4 mutant molecule and/or fragments thereof may be used to react with B7 positive cells, such as B cells, to regulate immune responses mediated by T cell interactions with the B7 antigen positive cells or in vitro for leukocyte typing so as to define B cell maturational stages and/or B cell associated diseases (Yokochi et al. J. Immuno. 128(2): 823). Surface immunostaining of leukocytes is accomplished by immunofluorescent technology or immunoenzymatic methods but other means of detection are possible.

Soluble CTLA4 mutant molecules, and/or fragments and derivatives of these proteins, may also be used to react with B7 positive cells, including B cells, to regulate immune responses mediated by T cell dependent B cell responses. The term "fragment" as used herein means a portion of the amino acid sequence encoding the protein referred to as "CTLA4" capable of binding B7. A fragment of the soluble CTLA4 mutant molecule that may be used is a polypeptide having an amino acid sequence corresponding to some portion of the amino acid sequence corresponding to the CTLA4 receptor used to obtain the soluble CTLA4 protein as described herein.

In one embodiment, CTLA4 mutant molecules may be introduced in a suitable pharmaceutical carrier in vivo, i.e. administered into a subject for treatment of pathological conditions such as immune system diseases or cancer (Pearson et al. "Transplantation Tolerance Induced by CTLA4-Ig" Transplantation, 1994 June 27, 57 (12):1701-6; Bolling et al. "The Effect of Combinations Cyclosporine and CTLA4-Ig Therapy on Cardiac Allograft Survival", Journal of Surgical Research, 1994 57(1):60-4).

Introduction of CTLA4 mutant molecules in vivo is expected to result in interference with T cell interactions with other cells, such as B cells, as a result of binding of the ligand to B7 positive cells. The prevention of normal T cell interactions may result in decreased T cell activity, for example, decreased T cell proliferation. In addition, administration of the mutant fusion protein in vivo is expected to result in regulation of in vivo levels of cytokines, including, but not limited to, interleukins, e.g. interleukin ("IL")-2, IL-3, IL-4, IL-6, IL-8, growth factors including tumor growth factor ("TGF"), colony stimulating factor ("CSF"), interferons ("IFNs"), and tumor necrosis factor ("TNF") to promote desired effects in a subject. For example, when the fusion protein is introduced in vivo, it may block production of cytokines, which contribute to malignant growth, for example of tumor cells. The fusion protein may also block proliferation of viruses dependent on T cell activation, such as the virus that causes AIDS, HTLV1.

Under some circumstances, as noted above, the effect of administration of the CTLA4 mutant molecule or its fragments in vivo is inhibitory, resulting from blocking by the fusion protein of the CTLA4 and CD28 triggering resulting from T cell/B cell contact. For example, the CTLA4 mutant molecule may block T cell proliferation. Introduction of the CTLA4 mutant molecule in vivo will thus produce effects on both T and B cell-mediated immune responses. The fusion protein may also be administered to a subject in combination with the introduction of cytokines or other therapeutic reagents.

In an additional embodiment of the invention, other reagents, including derivatives reactive with the CTLA4 mutant molecule are used to regulate T cell interactions. For example, antibodies, and/or antibody fragments reactive with the CTLA4 receptor may be screened to identify those capable of inhibiting the binding of the CTLA4 mutant molecule to the B7-1 antigen. The antibodies or antibody fragments such as Fab or F(ab)₂ fragments, may then be used to react

with the T cells, for example, to inhibit T cell proliferation.

In another embodiment, the CTLA4 mutant molecule may be used to identify additional compounds capable of regulating the interaction between CTLA4 or CD28 and the B7 antigens. Such compounds may include small naturally occurring molecules that can be used to react with B cells and/or T cells. For example, fermentation broths may be tested for the ability to inhibit CTLA4/B7 interactions. In addition, derivatives of the CTLA4lg mutant fusion protein as described above may be used to regulate T cell proliferation. For example, the fragments or derivatives may be used to block T cell proliferation in graft versus host (GVH) disease which accompanies allogeneic bone marrow transplantation.

The CD28-mediated T cell proliferation pathway is cyclosporine-resistant, in contrast to proliferation driven by the CD3/Ti cell receptor complex (June et al., *Mol. Cell. Biol.* 7:4472-4481 (1987)). Cyclosporine is relatively ineffective as a treatment for GVH disease (Storb, *Blood* 68:119-125 (1986)). GVH disease is thought to be mediated by T lymphocytes which express CD28 antigen (Storb and Thomas, *Immunol. Rev.* 88:215-238 (1985)). Thus, the CTLA4 mutant molecules may be useful alone, or in combination with immunosuppressants such as cyclosporine, for blocking T cell proliferation in GVH disease, and to treat other pathological conditions such as autoimmunity, transplantation rejection, infectious diseases and neoplasia.

The mutant CTLA4 molecules described herein may be formulated in a variety of dosage forms which include, but are not limited to, liquid solutions or suspensions, tablets, pills, powders, suppositories, polymeric microcapsules or microvesicles, liposomes, and injectable or infusible solutions. The preferred form depends upon the mode of administration and the therapeutic application.

The most effective mode of administration and dosage regimen for the molecules of the present invention depends upon the severity and course of the disease, the subject's health and response to treatment and the judgment of the treating physician. Accordingly, the dosages of the molecules should be titrated to the individual subject.

The interrelationship of dosages for animals of various sizes and species and humans based on mg/m² of surface area is described by Freireich, E.J., et al. (Quantitative Comparison of Toxicity of Anticancer Agents in Mouse, Rat, Hamster, Dog, Monkey and Man. *Cancer Chemother. Rep.*, 50, No.4, 219-244, May 1966).

Adjustments in the dosage regimen may be made to optimize the growth inhibiting response. Doses may be divided and administered on a daily basis or the dose may be reduced proportionally depending upon the situation. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the specific therapeutic situation.

In accordance with the practice of the invention an effective amount for treating a subject may be between about 0.1 and about 10mg/kg body weight of subject. Also, the effective amount may be an amount between about 1 and about 10 mg/kg body weight of subject.

ADVANTAGES OF THE INVENTION: CTLA4 mutant molecules of the invention are expected to be useful in vivo as an inhibitor of T cell activity, over non-specific inhibitors such as cyclosporine and glucocorticoids.

The B7-1 (CD80) antigen expressed on activated B cells and cells of other lineages, and the CTLA4 receptor expressed on T cells, can directly bind to each other, and this interaction can mediate cell-cell interaction. The B7-2 (CD86) antigen expressed on activated B cells and cells of other lineages, and the CD28 receptor expressed on T cells, can directly bind to each other, and this interaction can mediate cell-cell interaction. While B7-1 has a greater specificity for CTLA4 and B7-2 has a greater specificity for CD28, there is a certain amount of cross-reactivity between these ligands (Kuchroo et al., *Cell*, 80:707-718 (1995)).

Interactions between these ligands directly trigger the activation pathways in T cells, leading to cytokine production, T cell proliferation, and B cell differentiation into immunoglobulin producing cells. The activation of B cells that occurs, can cause increased expression of B7 antigen and further CD28 stimulation, leading to a state of chronic inflammation such as in autoimmune diseases, allograft rejection, graft versus host disease or chronic allergic reactions. Blocking or inhibiting this reaction may be effective in preventing T cell cytokine production and thus preventing or reversing inflammatory reactions.

Soluble CTLA4 molecules have been shown previously to be a potent inhibitor of in vitro lymphocyte functions requiring T and B cell interaction (see parent application, EP-A-0 682 039 and U.S. Serial No. 08/228,208). This indicates the importance of interactions between the B7 antigens and their counter-receptors, CTLA4 and/or CD28.

Recent data suggests that the B7-1 and B7-2 molecules differentially activate distinct T-cell subsets. Specifically, these antigens are linked to the differentiation of the CD4 T-helper cells, which upon stimulation, differentiate into two distinct subpopulations (designated Th1 and Th2), each producing its own set of cytokines and mediating separate effector functions. Significantly, these different subsets appear to play specific roles in a number of clinical pathologies (Cohen, *J. Science* 262, 175-176 (1993) and Simon et al, *P.N.A.S.* 91, 8562-8566 (1994)). By influencing the effector functions of these T-cell subsets, CTLA4 mutants molecules may prevent or reverse these types of pathologies.

Under conditions where T cell/B cell interactions are occurring as a result of contact between T cells and B cells, binding of introduced CTLA4 mutant molecules to react with B7 antigen positive cells, for example B cells, may interfere, i.e. inhibit, the T cell/B cell interactions resulting in regulation of immune responses. Because of this exclusively inhib-

itory effect, CTLA4 mutant molecules are expected to be useful in vivo as an inhibitor of T cell activity, over non-specific inhibitors such as cyclosporine and glucosteroids.

The following examples are presented to illustrate the present invention and to assist one of ordinary skill in making and using the same. The examples are not intended in any way to otherwise limit the scope of the invention.

EXAMPLE 1

By site-specific and homolog mutagenesis, we have identified regions in CTLA4Ig which are required for its high avidity binding to B7-1. The following is a description of how to make soluble CTLA4/CD28 hybrid fusion proteins which bind B7.

MATERIALS AND METHODS

Monoclonal antibodies (mAbs). Murine Mab's specific for CTLA4 were prepared and characterized as previously described (Linsley et al. J. Ex. Med., (1992) 176:1595-1604). Antibody 9.3 (anti-CD28) has been described previously ((Hansen et al., Immunogenetics 10:247-260 (1980)).

Cell Culture. The preparation of stably transfected B7-1 positive CHO cells has been previously described (Linsley et al., in J. Exp. Med. 173:721-730 (1991); P. S. Linsley et al., J. Exp. Med. 174:561 (1991)).

Cells were maintained in DMEM supplemented with 10% fetal bovine serum (FBS), 0.2mM proline, and 1μM methotrexate. COS cells were grown in DMEM supplemented with 10% FBS. CTLA4Ig was prepared in CHO cells as previously described (Example 2 of U.S. Serial No. 08/228,208, published with EP-A-0 682 039).

CTLA4Ig and CD28Ig site-directed mutant expression plasmids. Site-directed mutagenesis was performed on a vector encoding soluble chimeric form of CTLA4 (CTLA4Ig) in which the extracellular domain of CTLA4 was genetically fused to the hinge and constant regions of a human IgG heavy chain (Linsley et al, J. Exp. Med. 173:721-730 (1991)). CTLA4Ig site-directed mutants were prepared by encoding the desired mutation in overlapping oligonucleotide primers and generating the mutants by PCR (Ho et al., 1989, supra.) using the CTLA4Ig plasmid construct as a template.

Six mutants were prepared which encoded substitutions to alanine in the highly conserved hexapeptide 98MYPPPY103 forming part of the putative CDR3-like domain (Figure 1) (Ho et al., 1989, supra.). These mutants are described in Table II. In addition, two mutants encoding the residues P103A and Y104A (MYPPAY (SEQ ID NO:7) and MYPPPA (SEQ ID NO:8), respectively) from the CD28Ig 99MYPPPY104 hexapeptide using CD28Ig as a template were also prepared by the same method. These mutants are also described in Table II.

Primers required for PCR reactions but not for introducing mutations included (1) a CDM8 forward (CDM8FP) primer encoding a complementary sequence upstream of the HindIII restriction site at the 5' end of the CDM8 stuffer region, and (2) a reverse primer (CDM8RP) encoding a complementary sequence downstream of the XbaI site at the 3' end of the CDM8 stuffer region.

These primers encoded the following sequences:

CDM8FP:5'-AATACGACTCACTATAGG (SEQ ID NO:9)
CDM8RP:5'-CACCACACTGTATTAACC (SEQ ID NO:10)

PCR conditions consisted of 6 min at 94°C followed by 25 cycles of 1 min at 94°C, 2 min at 55°C and 3 min at 72°C. Taq polymerase and reaction conditions were used as suggested by the vendor (Perkin Elmer Cetus, Emeryville, CA). PCR products were digested with HindIII and XbaI and ligated to HindIII/XbaI-cut CDM8 expression vector.

To confirm that the desired mutations had been inserted and to verify the absence of secondary mutations, each CTLA4Ig mutant fusion protein (an example of a soluble CTLA4 mutant fusion protein) was sequenced by the dideoxy chain termination/extension reaction with Sequenase reagents used according to the manufacturer's recommendations (United States Biochemical Corp., Cleveland, OH).

Plasmids were transfected into COS cells (Aruffo et al., Cell 61:1303 (1990)) and the conditioned media was used as a source for the resulting Ig mutant fusion proteins.

CTLA4/CD28Ig hybrid expression plasmids. CTLA4/CD28Ig hybrid scan plasmids encoding the constructs HS2, HS4, HS4-A, HS4-B, and HS5 (Figure 3 and Table I) were prepared by PCR using overlapping oligonucleotide primers designed to introduce CTLA4 sequences into CD28Ig while, at the same time, deleting the equivalent region from CD28. The same CDM8 forward and reverse PCR primers described above were also used.

The following is a list of the CTLA4/CD28 hybrid fusion proteins which were made.

DESIGNATION	FRAMEWORK	MODIFICATIONS
HS1	CTLA4	1-24 OF CD28

(continued)

5

10

15

20

25

30

DESIGNATION	FRAMEWORK	MODIFICATIONS
HS2	CD28	97-125 OF CD28 1-22 OF CTLA4 96-125 OF CTLA4
HS3	CTLA4	96-125 OF CD28
HS4	CD28	96-123 OF CTLA4
HS4A	CD28	96-113 OF CTLA4
HS4B	CD28	114-123 OF CTLA4
HS5	CD28	25-32 OF CTLA4
HS6	CTLA4	25-32 OF CD28
HS7	CD28	96-123 OF CTLA4 25-32 OF CTLA4
HS8	CD28	25-32 OF CTLA4 96-113 OF CTLA4
HS9	CD28	25-32 OF CTLA4 114-123 OF CTLA4
HS10	CD28	96-123 OF CTLA4 51-58 OF CTLA4
HS11	CD28	25-32 OF CTLA4 51-58 OF CTLA4
HS12	CD28	96-123 OF CTLA4 51-58 OF CTLA4 96-113 OF CTLA4
HS13	CD28	25-32 OF CTLA4 51-58 OF CTLA4 96-113 OF CTLA4
HS14	CD28	51-58 OF CTLA4

35

Each cDNA construct was genetically linked to cDNA encoding the hinge and constant regions of a human IgG1 in order to make soluble chimeras.

A HS6 hybrid was prepared in a similar manner to that described above except that the CDR1-like region in CTLA4Ig was replaced with the equivalent region from CD28Ig.

40

HS7, HS8, and HS9 constructs were prepared by replacing a ≈350 base-pair HindIII/HpaI 5' fragment of HS4, HS4-A, and HS4-B, respectively, with the equivalent cDNA fragment similarly digested from HS5 thus introducing the CDR1-like loop of CTLA4 into those hybrids already containing the CTLA4 CDR3-like region.

45

HS10-HS13 constructs are domain homolog mutants which were prepared by introducing the CDR2-like loop of CTLA4Ig into previously constructed homolog mutants. This was done by overlapping PCR mutagenesis whereby primers were designed to introduce CTLA4 CDR2-like sequences into homolog templates while at the same time deleting the equivalent CD28 CDR2-like region from the molecule.

Accordingly, HS4 served as a template to make HS10; HS7 served as a template to make HS11; HS4-A served as a template to make HS12; and HS8 served as a template to make HS13 (Figure 3 and Table I). The CDM8 primers described above were also used in these constructions.

50

The HS14 hybrid construct was prepared by replacing the CDR2-like loop of CD28 with the equivalent loop from CTLA4Ig (Figure 3 and Table I).

Oligonucleotide primers designed to introduce these changes were used in overlapping PCR mutagenesis identical to that described for other mutants.

PCR reactions and subcloning into CDM8 were performed as described above. Again all mutants were sequenced by the dideoxy chain termination/extension reaction.

55

Plasmids encoding each of the mutants were transfected into COS cells and the resulting soluble Ig fusion proteins were quantitated in culture media and visualized by Western blot as described in following sections.

Quantitation of the resulting Ig fusion proteins in culture media. Soluble mutant fusion proteins were quantitated

in an enzyme immunoassay by determining the amount of Ig present in serum-free COS cell culture media.

Microtiter plates (Immulon2; Dynatech Labs., Chantilly, VA) were coated with 0.5µg/ml goat anti-human IgG (Jackson ImmunoResearch Labs., West Chester, PA) for 16-24h at 4°C. Wells were blocked for 1h with specimen diluent (Genetic Systems, Seattle, WA), then washed with PBS containing 0.05% Tween 20 (PBS-Tw).

COS cell culture media containing fusion proteins was added at various dilutions and incubated for 1h at 22°C. Known concentrations of CTLA4Ig were also added to separate wells on each plate for a standard curve.

After washing, horseradish peroxidase (HRP)-conjugated goat anti-human IgG (Tago, Burlingame, CA) diluted 1:12,000 was added and incubated for 1h at 22°C. Wells were then washed and incubated with 3,3',5,5' tetramethylbenzidine (TMB) substrate (Genetic Systems) for 15 min before stopping the reaction by the addition of 1N H₂SO₄. Optical density was measured at dual wavelengths of 450 and 630nm on a microtiter plate reader (Genetic Systems).

Concentration of mutant Ig fusion protein was determined by comparison with a standard curve of known concentrations of CTLA4Ig.

Immunoprecipitation and Western blot analysis. CTLA4/CD28Ig hybrid fusion proteins present in culture media were adsorbed to protein A-Sepharose by overnight incubation at 4°C. The beads were washed with PBS containing 0.1% Nonidet-P40 (NP40) then SDS PAGE sample buffer was added and the eluted protein was loaded onto an SDS polyacrylamide gel.

Western blot transfer of protein onto nitrocellulose was done by standard procedures. Nitrocellulose membranes were then blocked with PBS containing 0.1% NP40 and 1% non-fat dry milk powder.

After washing in PBS-Tw membranes were incubated with alkaline phosphatase-conjugated goat anti-human IgG (Boehringer Mannheim, Indianapolis, IN) diluted 1:1,000 and incubated for 1h at 22°C. Blots were then washed and developed using standard procedures.

B7 positive CHO cell enzyme immunoassay. The ability of CTLA4Ig mutant fusion proteins, and CTLA4/CD28Ig hybrid fusion proteins to bind B7-1 stably expressed on CHO cells was determined by an enzyme immunoassay.

Round bottom tissue culture treated 96 well microtiter plates (Corning, Corning, NY) were seeded with B7-1 positive CHO cells at 10³ cells/well. Two days later the confluent cells were fixed in 95% ethanol for 15 min.

After washing with PBS-Tw, mutant Ig fusion proteins were added at various concentrations and incubated for 1h at 4°C. After washing, HRP-conjugated goat anti-human IgG (Tago) diluted 1:10,000 was added and incubated for 1h at 22°C.

Wells were then washed and TMB substrate added as above and allowed to react for 30 min before stopping the reaction with 1N H₂SO₄.

Absorbance of the wells was measured at 450nm.

CD28Ig site-directed mutant fusion protein binding assay. Site-directed mutant fusion proteins of CD28Ig were assayed for their ability to bind to B7-1 by an indirect enzyme immunoassay.

Wells of ELISA plates were coated with a chimeric fusion protein containing the extracellular domain of human B7-1 fused to a mouse IgG1 Fc region, at 5µg/ml for 16h at 4°C. Wells were blocked for 1h with specimen diluent (Genetic Systems) then washed with PBS-Tw. COS cell culture media containing known concentrations of mutant fusion protein was added at various concentrations and incubated for 1h at 22°C.

Known concentrations of CD28Ig were also added to separate wells on each plate. After washing, HRP-conjugated goat anti-human IgG (Tago) diluted 1:10,000 was added and incubated for 1h at 22°C. TMB substrate was added and optical densities read as described for quantitation of Ig fusion proteins in culture media.

mAb binding to Ig fusion proteins. The ability of anti-CTLA4 mAb's and the anti-CD28 mAb 9.3 to bind CTLA4/CD28Ig hybrid fusion proteins and CTLA4Ig mutant fusion proteins was assessed by an enzyme immunoassay.

Wells of microtiter plates (Immulon 2) were coated with 0.5µg/ml of goat anti-human IgG (Jackson) for 16-24h at 4°C. Plates were blocked for 1h with specimen diluent (Genetic Systems), washed with PBS-Tw, then incubated with the Ig fusion proteins for 1h at 22°C. After washing, wells were incubated with mAb at 1µg/ml for 1h at 22°C.

After further washing, HRP-conjugated goat anti-mouse Ig (Tago) diluted 1:10,000 was added and incubated for 1h at 22°C. TMB substrate was added and optical density measured as described above.

CTLA4 molecular model. An approximate three-dimensional model of the CTLA4 extracellular domain was generated based on the conservation of consensus residues of IGSF variable-like domains.

Using such IGSF consensus residues as "anchor points" for sequence alignments, CTLA4 residues were assigned to the A, B, C, C', C'', D, E, F, G strands of an Ig variable fold (Williams/Barclay, 1988, supra.) and the connecting loop regions (Figure 6).

The CTLA4 model was built (InsightII, Discover, Molecular Modeling and Mechanics Programs, respectively, Biosym Technologies, Inc., San Diego) using the variable heavy chain of HyHEL-5 (Sheriff et al., 1987 PNAS 84: 8075-8079) as template structure. Side-chain replacements and loop conformations were approximated using conformational searching (Brucoleri et al., 1988 335:564-568).

Several versions of the model with modified assignments of some residues to β-strands or loops were tested using 3D-profile analysis (Lüthy et al., 1992, Nature 336:83-85) in order to improve the initial alignment of the CTLA4 extra-

cellular region sequence with an IGSF variable fold.

RESULTS

Construction and binding activity of CTLA4Ig and CD28Ig mutant fusion proteins. A sequence alignment of various homologues of CD28 and CTLA4 is demonstrated in Figure 1. In Figure 1, sequences of human (H), mouse (M), rat (R), and chicken (Ch) CD28 are aligned with human and mouse CTLA4. Residues are numbered from the mature protein N-terminus with the signal peptides and transmembrane domains underlined and the CDR-analogous regions noted. Dark shaded areas highlight complete conservation of residues while light shaded areas highlight conservative amino acid substitutions in all family members.

Regions of sequence conservation are scattered throughout the extracellular domains of these proteins with the most rigorous conservation seen in the hexapeptide MYPPPY (SEQ ID NO:11) motif located in the CDR3-like loop of both CTLA4 and CD28 (Figure 1). This suggests a probable role for this region in the interaction with a B7 antigen, e.g., B7-1 and B7-2.

To test this possibility, site-directed alanine scanning mutations were introduced into this region of CTLA4Ig using PCR oligonucleotide primer-directed mutagenesis thereby resulting in CTLA4Ig mutant fusion proteins. Similarly two alanine mutations were introduced into the CD28Ig MYPPPY motif thereby resulting in CD28Ig mutant fusion proteins.

All cDNA constructs were sequenced to confirm the desired mutations before transfection into COS cells. The concentrations of mutant Ig fusion proteins in serum-free COS cell culture media were determined by an Ig quantitation assay.

The ability of each CTLA4Ig mutant fusion protein to bind to B7-1 expressed on stably transfected CHO cells was then determined by an indirect cell binding immunoassay. Binding of CD28Ig mutant fusion proteins to B7-1 was assessed by an indirect enzyme immunoassay. Each of these assays are described in Materials and Methods.

Mutagenesis of each residue of the CTLA4Ig MYPPPY motif to Ala had a profound effect on binding to B7-1 as shown in Figure 2. Figure 2 shows that mutations in the MYPPPY motif of CTLA4Ig and CD28Ig disrupt binding to B7-1. Site-directed mutant Ig fusion proteins were produced in transiently transfected COS cells, quantitated and tested for their ability to bind to B7-1.

In Figure 2 fusion protein quantitations were repeated at least twice with replicate determinations. Specifically, Figure 2 shows that CTLA4Ig mutants bind to stably transfected, ethanol-fixed B7-1+ CHO cells grown to confluency in ELISA tissue culture plates. Binding data is expressed as the average of duplicate wells and is representative of at least two experiments.

Y99A and P101A mutants bound to B7-1 but with considerably reduced ability relative to wild-type CTLA4Ig. In contrast, the mutants M98A, P100A, P102A and Y103A showed an almost complete loss of binding. Furthermore, the CD28Ig MYPPPY mutants P103A and Y104A did not display detectable binding to B7-1 immobilized on wells of ELISA plates (Figure 2).

B7-1 transfected CHO cells which were incubated with CTLA4Ig mutant fusion protein, labeled with anti-human FITC, and assayed using a FACSCAN showed equivalent results. These results clearly demonstrate a critical role for the MYPPPY motif in both CTLA4Ig and CD28Ig binding to B7-1.

Characterization of CTLA4/CD28Ig hybrid fusion proteins. Since the MYPPPY motif is common to both CTLA4Ig and CD28Ig, it alone cannot account for the observed differences in binding to B7-1 seen with CTLA4Ig and CD28Ig. The contribution of less well conserved residues to high avidity binding B7-1 was assessed using a series of homolog mutants.

The three CDR-like regions of CD28 were replaced in various combinations with the equivalent regions from the CTLA4 extracellular domain (Figure 3 and Table I). Figure 3 is a map of CTLA4/CD28Ig mutant fusion proteins showing % binding activity to B7-1+ CHO cells relative to CTLA4Ig. Conserved cysteine residues (C) are shown at positions 22, 93 and 121 respectively (CTLA4 numbering). Also shown is the position of the MYPPPY motif. Open areas represent CD28 sequence; filled areas represent CTLA4 sequence; cross-hatched areas represent beginning of IgG Fc (also refer to Table I). Percent binding activities were determined by comparing binding curves (Figures 4/5) relative to CTLA4Ig and finding the concentration of a mutant required to give the same O.D. as that found for CTLA4Ig. The ratio of mutant protein to CTLA4Ig concentration at a particular O.D. was then expressed as % binding activity. At least two A450 readings were taken from the linear part of the CTLA4Ig binding curve and the average % binding activity determined.

A total of 14 hybrid cDNA constructs were prepared, sequenced, and transfected into COS cells. Concentrations of Ig fusion proteins in serum-free culture media were determined and their electrophoretic mobility compared by SDS-PAGE including Western blotting analysis.

Under reducing conditions each chimeric protein migrated with a relative molecular mass ranging between that of CTLA4Ig (Mr-50kDa) and CD28Ig (Mr-70kDa) depending on the size of the exchanged region.

Under non-reducing conditions the proteins migrated primarily between 100-140kDa indicating that these fusion

proteins existed as disulfide-linked dimers despite mutagenesis of the cysteine residues in the hinge region of the Fc.

Since four of the five conserved cysteine residues in CTLA4 and CD28 are thought to be involved in intrachain disulfide bonds, dimerization of the fusion proteins was therefore most likely attributable to the fifth conserved cysteine residue at position 121 in CTLA4 (position 123 in CD28).

Binding of CTLA4/CD28Ig hybrid fusion proteins to B7-1. The hybrid fusion proteins were tested for their ability to bind to B7-1 by the same indirect cell binding immunoassay used to assay the site-specific CTLA4Ig and CD28Ig mutant fusion proteins.

Under these conditions the binding between CD28Ig and B7-1 is barely detectable (Figures 4/5). However, replacing residues 97 to 125 (the CDR3-like extended region) of CD28 with the corresponding residues of CTLA4 resulted in an approximately two and a half orders of magnitude increase in binding of the CD28Ig analog to B7-1 (Figures 4/5). Figures 4/5 shows that CTLA4/CD28Ig mutant fusion proteins demonstrate involvement of CDR-analogous regions in high avidity binding to B7-1 CHO cells. Mutants were assayed as described in figure 2. Data is expressed as the average of duplicate wells and is representative of at least three experiments. From these curves % binding activity relative to CTLA4Ig was determined as explained and shown in Figure 3.

Binding to B7-1 by this construct, termed HS4 (Figure 3), is approximately five fold less than wild type CTLA4Ig. The HS2 hybrid which includes additional N-terminal residues of CTLA4 (amino acids 1-22), did not improve the ability of the hybrid molecule to bind to B7-1 relative to HS4.

The HS6 construct which represents the CTLA4Ig sequence except that it contains the CDR1-like region of CD28 (residues 25-32), bound similarly. However, the additional inclusion of the CTLA4 CDR1-like region (residues 25-32) into the HS4 construct (termed HS7), showed further improved binding so that the binding affinity is approximately 44% of CTLA4Ig (Figure 3).

In contrast, inclusion of the CDR2-like region of CTLA4 (residues 51-58) into HS4 (construct HS10), did not further increase binding (Figure 3). A similar result was found for construct HS11 which had all three CDR-like region sequences of CTLA4 included into CD28Ig. The HS5 hybrid which contained only the CDR1-like domain of CTLA4 bound at very low levels.

The CTLA4/CD28Ig hybrid HS4-A encoded CTLA4Ig residues 96-113 in the C-terminally extended CDR3-like region; nine CTLA4 derived residues fewer than HS4 (Figure 3 and Table I). HS4-A bound B7-1 CHO cells less well than HS4 (Figures 3 and 5). However, addition of the CTLA4 CDR1-like loop (HS8 hybrid), increased B7-1 binding from about 2% to nearly 60% of wild type binding.

On the other hand, addition of the CTLA4 CDR2-like loop into HS4-A (HS12) did not increase binding relative to HS4-A; neither did addition of all three CTLA4 CDR-like regions (HS13, Figure 3).

Another hybrid called HS4-B, encoded the CD28 CDR3-like region including the MYPPPY motif followed by CTLA4 residues 114-122 (Table I and Figure 3).

HS4-B and HS4-A displayed similar binding to B7-1. Unlike HS4-A, however, the inclusion of the CTLA4 CDR1-like loop into HS4-B (HS9) did not improve binding (Figure 3), suggesting that residues immediately adjacent to the CTLA4Ig MYPPPY motif were important determinants in high avidity binding.

Monoclonal antibody binding to CTLA4/CD28Ig hybrid fusion proteins. The structural integrity of each hybrid fusion protein was examined by assessing their ability to bind mAb's specific for CTLA4 or CD28 in an enzyme immunoassay. The CTLA4 specific mAb's 7F8, 11D4 and 10A8 block ligand binding (Linsley et al. (1992) supra.).

These antibodies bound to each of the CTLA4Ig mutant fusion proteins except 11D4 which failed to bind to P100A and P102A (Table II). Since 7F8 and 10A8 bound to these mutants, the lack of binding by 11D4 can probably be attributed to mutagenesis perturbing the epitope recognized by 11D4.

Conversely, each antibody failed to bind to any of the homolog scan hybrid fusion proteins except 7F8 which bound to HS6, and 11D4 which bound weakly to HS8. As many of these homolog hybrid fusion proteins were, to some extent, able to bind to B7-1, it is likely that lack of binding by the antibodies was due to disruption of conformational epitopes formed by spatially adjacent but non-linear sequences.

The CD28 specific mAb 9.3 (Linsley et al. (1992) supra.) failed to bind to either of the CD28 site-directed mutant fusion proteins but bound to the hybrid fusion proteins HS4, HS4-A, HS7 and HS8. With HS2, weaker binding was observed. No binding was seen with the HS5 and HS6 constructs.

CTLA4 model. Figure 6 shows a schematic representation of the CTLA4 model. The assignment of CTLA4 residues to CDR-like regions is shown in Figure 1. The CTLA4 model suggests the presence of an additional (non-Ig) disulfide bond between residues Cys49 and Cys67 which supports the similarity of CTLA4 and the Ig variable fold.

The two possible N-linked glycosylation sites in CTLA4 map to solvent exposed positions of the Ig β -strand framework regions. 3D-profile analysis indicated that the CTLA4 sequence is overall compatible with an Ig V-fold, albeit more distantly related.

Residue Val115 represents the last residue of the CTLA4Ig-like domain. The conformation of the region between Val115 and the membrane-proximal Cys121 which is thought to form the CTLA4 homodimer is highly variable in the CD28 family. The picture that emerges is that CD28 family members mainly utilize residues in two of three CDR-like

regions for binding to B7-1.

The MYPPPY motif represents a conserved scaffold for binding which appears to be augmented by its C-terminal extension and which is specifically modulated by the highly variable CDR1-like region. CDR3 and CDR1-like regions are spatially contiguous in Ig-variable folds. The CDR2 like region is spatially distant and does not, in the case of the CD28 family, significantly contribute to the binding to B7-1.

EXAMPLE 2

By site-specific mutagenesis, we have identified an amino acid position in CTLA4 which plays a crucial role in this molecule's ability to bind both CD80 and CD86. In addition we identify 2 amino acid substitutions at this position which generate mutant CTLA4Ig molecules that have the ability to bind CD80 in a manner similar to the wild type molecule, but have lost the ability to bind CD86. The following is a description of how to make soluble CTLA4 fusion proteins which bind CD80 but not CD86.

MATERIALS AND METHODS

Monoclonal antibodies (mAbs). Murine monoclonal antibodies specific for CD80 and CD86 have been described previously (Kuchroo et al., *Cell*, Vol. 80, 707-718, (1995)).

Cell Culture. The preparation of stably transfected B7-1 (CD80) positive CHO cells has been previously described (Linsley et al., in *J. Exp. Med.* 173:721-730 (1991); P. S. Linsley et al., *J. Exp. Med.* 174:561 (1991)). The preparation of stably transfected B7-2 (CD86) cells has been previously described (Freeman et al., *J. Exp. Med.* 178: 2185 (1993); PCT WO95/06738). The preparation of LCL cells has been previously described (Wyss-Coray et al., *European Journal of Immunology*, 23(12), 3350-3357 (1993)).

Cells were maintained in DMEM supplemented with 10% fetal bovine serum (FBS), 0.2mM proline, and 1μM methotrexate. COS cells were grown in DMEM supplemented with 10% FBS. CTLA4Ig was prepared in CHO cells as previously described (see parent application, U.S. Serial No. 08/228,208, Example 2, and EP-A-0 682 039).

CTLA4Ig site-directed mutant expression plasmids. Site-directed mutagenesis was performed on a vector encoding soluble chimeric form of CTLA4 (CTLA4Ig) in which the-extracellular domain of CTLA4 was genetically fused to the hinge and constant regions of a human IgG heavy chain (Linsley et al., *J. Exp. Med.* 173:721-730 (1991)). CTLA4Ig site-directed mutants were prepared by encoding the desired mutation in overlapping oligonucleotide primers and generating the mutants by PCR (Ho et al., 1989, *supra*.) using the πLN CTLA4Ig plasmid construct as a template (see parent application, U.S. Serial No. 08/228,208, Example 1, and EP-A-0 682 039).

Eighteen mutants were prepared which encode amino acid substitutions at the first tyrosine position in the highly conserved MYPPPY motif of CTLA4 (Figure 1) (Ho et al., 1989, *supra*). This series of mutants was generated by manipulating the codon at this position in order to generate new codons encoding each of the twenty amino acids except cysteine.

Primers required for PCR reactions but not for introducing mutations included (1) a πLN forward primer(πLNlgFP) encoding a complementary sequence upstream of the SacI restriction site at the 5' end of the CTLA4Ig πLN sequence, and (2) a reverse primer (πLNlgRP) encoding a complementary sequence downstream of the XbaI site at the 3' end of the immunoglobulin encoding region.

These primers encoded the following sequences:

πLNlgFP:5'-TGCAAGGTGGAGCTCATGTTCCACCGCCATAC (SEQ ID NO:12)

πLNlgRP:5'-GCGCTCGACTCTAGAAGCATCCTCGTG (SEQ ID NO:13)

PCR conditions consisted of 6 min at 94°C followed by 30 cycles of 1 min at 94°C, 2 min at 55°C and 3 min at 72°C. Taq polymerase and reaction conditions were used as suggested by the vendor (Perkin Elmer Cetus, Emeryville, CA). PCR products were digested with SacI and XbaI and ligated to SacI/XbaI-cut πLN CTLA4Ig expression vector.

To confirm that the desired mutations had been inserted and to verify the absence of secondary mutations, each CTLA4Ig mutant fusion protein (an example of a soluble CTLA4 mutant fusion protein) was sequenced by the dideoxy chain termination/extension reaction with Sequenase reagents used according to the manufacturers recommendations (United States Biochemical Corp., Cleveland, OH).

Plasmids encoding each of the mutants were transfected into COS cells (Aruffo et al., *Cell* 61:1303 (1990)) and the conditioned media was used as a source for the resulting Ig mutant fusion proteins.

Quantitation of the resulting Ig fusion proteins in culture media. Soluble mutant fusion proteins were quantitated by FACS analysis as previously described (see parent application, U.S. Serial No. 08/228,208, Example 1, and EP-A-0 682 039).

Binding CTLA4Ig fusion proteins in the presence of anti-CD80 or anti-CD86 mAbs. Binding studies involving

CTLA4Ig proteins and competitor molecules utilized previously described procedures (see parent application, U.S. Serial No. 08/228,208, Example 4, and EP-A-0 682 039).

Purification of Ig fusion proteins. Ig fusion proteins were purified as previously described (See parent application, U.S. Serial No. 08/228,208, Example 1, and EP-A-0 682 039).

RESULTS

Binding activity of CTLA4Ig mutant fusion proteins. Site-directed mutant Ig fusion proteins were produced in transiently transfected COS cells, quantitated and tested for their ability to bind to CD80 or CD86.

The ability of each CTLA4Ig mutant fusion protein to bind CD80 or CD86 expressed on stably transfected CHO cells was determined by incubating these cells with the different CTLA4Ig mutant fusion proteins, labelling the mutant proteins with anti-human FITC, and assaying them via FACSCAN as described above.

Titration staining of CHO cells which express either CD80 or CD86 shows that mutagenesis of the first tyrosine residue in the CTLA4Ig MYPPPY motif has a profound effect on binding to both CD80 and CD86. Figure 7 shows that each amino acid substitution at this position produces a unique binding profile. In addition, this figure shows that substitutions with phenylalanine and tryptophan at this position generate mutants that are able to bind CD80 in a manner analogous to the wild type molecule, but are unable to bind CD86.

Figure 8 further illustrates the unique characteristics of the two mutant molecules in which the first tyrosine in the MYPPPY motif has been replaced with either phenylalanine or tryptophan. FACS analysis over a range of different concentrations shows that these two mutants bind CD80 in a manner analogous to the wild type molecule but are completely unable to bind to CD86.

These observations are consistent with studies involving cell lines which naturally express the CD80 and CD86 antigens. Figure 9 shows that the phenylalanine mutant binds to endogenous CD80 which is expressed on the surface of the cell line LCL 816. This mutant's specificity for the CD80 molecule is demonstrated in competition studies where its binding is inhibited by monoclonal antibodies which are specific for CD80. Further, antibodies specific for CD86 have no effect on this molecule's ability to bind to this cell line.

This data clearly demonstrates that the first tyrosine in the MYPPPY motif in CTLA4Ig plays a critical role in this molecule's ability to bind both CD80 and CD86. In addition, these results show that substitutions which change the residue at this position to either a phenylalanine or a tryptophan generate mutants which retain the ability to bind CD80 but lose the ability to bind CD86.

TABLE I. CTLA4/CD28Ig homolog mutant junction sequences.

MUTANT

HS1	-22CKYasp27-	-93ckvEVM99-	-123CPSDQE-
HS2	-20fvcKYS25-	-94CKIelm98-	-121cpdDQE-
HS3		-93ckvEVM99-	-123CPSDQE-
HS4		-94CKIelm98-	-121cpdDQE-
HS5	-22CKYasp27-		-123CPSDQE-
HS6	-22ceySYN27-		-121cpdDQE-
HS4-A		-30ateFRA35-	-123CPSDQE-
HS4-B		-30SREvr35-	-121cpdDQE-
HS7	-22CKYasp27-		-123CPSDQE-
HS8	-22CKYasp27-		-121cpdDQE-
HS9	-22CKYasp27-		-123CPSDQE-
HS10		-30ateFRA35-	-121cpdDQE-
HS11	-22CKYasp27-		-123CPSDQE-
HS12		-30ateFRA35-	-121cpdDQE-
HS13	-22CKYasp27-		-123CPSDQE-
HS14		-30ateFRA35-	-121cpdDQE-

Junction sequences of the CTLA4/CD28-Ig hybrid fusion proteins. Amino acids are denoted by their single letter code with those in upper case being CD28 residues, those in lower case being CTLA4 residues and those in bold upper case being human IgG1 residues. Numbering in the table is from the N-terminal methionine of the respective proteins and refers to the adjacent amino acids.

TABLE II.

Binding of CTLA4 and CD28 monoclonal antibodies to CTLA4Ig and CD28Ig mutant fusion proteins and to CTLA4/CD28Ig hybrid fusion proteins.

	anti-CTLA4 mAbs			anti-CD28 mAb
	7F8	11D4	10A8	9.3
CTLA4Ig MUTANT FUSION PROTEIN				
AYPPPY	+++	+++	+++	-
MAPPPY	++	+	++	-
MYAPPY	+	-	+	-
MYPAPY	+++	+++	+++	-
MYPPAY	+++	-	+	-
MYPPPA	+++	++	+++	-
AAPPPY	+	++	+++	-
CD28Ig MUTANT FUSION PROTEIN				
MYPPAY	-	-	-	-
MYPPPA	-	-	-	+
CTLA4/CD28Ig HYBRID FUSION PROTEINS				
HS1	-	-	-	-
HS2	-	-	-	+
HS3	-	-	-	-
HS4	-	-	-	+++
HS5	-	-	-	-
HS6	+	-	-	-
HS4-A	-	-	-	++
HS4-B	-	-	-	++
HS7	-	-	-	+++
HS8	-	+	-	+++
HS9	-	+	-	-
HS10	-	-	-	-
HS11	-	-	-	+
HS12	-	-	-	-
HS13	-	-	-	-
HS14	-	-	-	-
CTLA4Ig	+++	+++	+++	-
CD28Ig	-	-	-	+++

Antibody binding was rated from that seen for wild type protein (+++) to above background (+), and no detectable binding (-).

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: BRISTOL-MYERS SQUIBB COMPANY
 (B) STREET: 345 Park Avenue
 (C) CITY: New York
 (D) STATE: New York
 (E) COUNTRY: USA
 (F) POSTAL CODE (ZIP): NY 10154

(ii) TITLE OF INVENTION: CTLA4 mutant molecules and uses thereof

(iii) NUMBER OF SEQUENCES: 13

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
 (B) COMPUTER: IBM PC compatible
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: EP 96305298.0

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 234 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Met	Ala	Cys	Leu	Gly	Phe	Gln	Arg	His	Lys	Ala	Gln	Leu	Asn	Leu	Ala	1	5	10	15
Ala	Arg	Thr	Trp	Pro	Cys	Thr	Leu	Leu	Phe	Phe	Leu	Leu	Phe	Ile	Pro	20	25	30	
Val	Phe	Cys	Lys	Ala	Met	His	Val	Ala	Gln	Pro	Ala	Val	Val	Leu	Ala	35	40	45	
Ser	Ser	Arg	Gly	Ile	Ala	Ser	Phe	Val	Cys	Glu	Tyr	Ala	Ser	Pro	Gly	50	55	60	
Lys	Ala	Thr	Glu	Val	Arg	Val	Thr	Val	Leu	Arg	Gln	Ala	Asp	Ser	Gln	65	70	75	80
Val	Thr	Glu	Val	Cys	Ala	Ala	Thr	Tyr	Met	Thr	Gly	Asn	Glu	Leu	Thr	85	90	95	
Phe	Leu	Asp	Asp	Ser	Xaa	Xaa	Ile	Cys	Thr	Gly	Thr	Ser	Ser	Gly	Asn	100	105	110	
Gln	Val	Asn	Leu	Thr	Ile	Gln	Gly	Leu	Arg	Ala	Met	Asp	Thr	Gly	Leu	115	120	125	
Tyr	Ile	Cys	Lys	Val	Glu	Leu	Met	Tyr	Pro	Pro	Pro	Tyr	Tyr	Leu	Xaa	130	135	140	
Gly	Ile	Gly	Asn	Gly	Thr	Gln	Ile	Tyr	Val	Ile	Asp	Pro	Glu	Pro	Cys	145	150	155	160

Xaa	Xaa	Xaa	Xaa	Xaa 165	Pro	Asp	Ser	Asp	Phe 170	Leu	Leu	Trp	Ile	Leu 175	Ala
Ala	Val	Ser	Ser 180	Gly	Leu	Phe	Phe 185	Tyr	Ser	Phe	Leu	Leu	Thr 190	Xaa	Ala
Val	Ser	Leu 195	Ser	Lys	Met	Leu	Lys 200	Lys	Arg	Ser	Pro	Leu 205	Thr	Thr	Gly
Val 210	Tyr	Val	Lys	Met	Pro	Pro 215	Thr	Glu	Pro	Glu	Cys 220	Glu	Xaa	Xaa	Lys
Gln 225	Phe	Gln	Pro	Tyr	Phe 230	Ile	Pro	Ile	Asn						

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 234 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met 1	Ala	Cys	Leu 5	Gly	Leu	Arg	Arg	Tyr	Lys 10	Ala	Gln	Leu	Gln	Leu 15	Pro
Ser	Arg	Thr	Trp 20	Pro	Phe	Val	Ala	Leu 25	Leu	Thr	Leu	Leu	Phe 30	Ile	Pro
Val	Phe	Ser 35	Glu	Ala	Ile	Gln 40	Val	Thr	Gln	Pro	Ser	Val 45	Tyr	Leu	Ala
Ser	Ser 50	His	Gly	Tyr	Ala	Ser 55	Phe	Pro	Cys	Glu	Tyr 60	Ser	Pro	Ser	His
Asn 65	Thr	Asp	Glu	Tyr	Arg 70	Val	Thr	Val	Leu	Arg 75	Gln	Thr	Asn	Asp	Gln 80
Met	Thr	Glu	Val	Cys 85	Ala	Thr	Thr	Phe	Thr 90	Glu	Lys	Asn	Thr	Val 95	Gly
Phe	Leu	Asp	Tyr 100	Pro	Xaa	Xaa	Phe	Cys 105	Ser	Gly	Thr	Phe	Asn 110	Glu	Ser
Arg	Val 115	Asn	Leu	Thr	Ile	Gln 120	Gly	Leu	Arg	Ala	Val	Asp 125	Thr	Gly	Leu

~~Tyr Leu Cys Lys Val Glu Leu Met Tyr Pro Pro Pro Tyr Phe Val Xaa~~
130 135 140

Gly Met Gly Asn Gly Thr Gln Ile Tyr Tyr Ile Asp Pro Glu Pro Cys
145 150 155 160

Xaa Xaa Xaa Xaa Xaa Pro Asp Ser Asp Phe Leu Leu Trp Ile Leu Tyr
165 170 175

Ala Val Ser Leu Gly Leu Phe Phe Tyr Ser Phe Leu Val Ser Xaa Ala
180 185 190

Val Ser Leu Ser Lys Met Leu Lys Lys Arg Ser Pro Leu Thr Thr Gly
195 200 205

Val Tyr Val Lys Met Pro Pro Thr Glu Pro Glu Cys Glu Xaa Xaa Lys
210 215 220

Gln Phe Gln Pro Tyr Phe Ile Pro Ile Asn
225 230

5

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 225 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

15

Met Thr Leu Arg Leu Leu Phe Leu Ala Leu Asn Phe Phe Xaa Ser Val
1 5 10 15

Gln Val Thr Glu Asn Lys Ile Leu Val Lys Gln Ser Pro Leu Leu Tyr
20 25 30

20

Val Asp Ser Asn Glu Val Xaa Ser Leu Ser Cys Arg Tyr Ser Tyr Asn
35 40 45

Leu Leu Ala Lys Glu Phe Arg Ala Ser Leu Tyr Lys Gly Val Asn Ser
50 55 60

25

Asp Val Xaa Glu Val Cys Val Gly Asn Gly Asn Phe Thr Tyr Gln Pro
65 70 75 80

Gln Phe Arg Ser Asn Ala Glu Phe Asn Cys Asp Gly Asp Phe Asp Asn
85 90 95

30

Glu Thr Val Thr Phe Arg Leu Trp Asn Leu His Val Asn His Thr Asp
100 105 110

Ile Tyr Phe Cys Lys Ile Glu Phe Met Tyr Pro Pro Pro Tyr Leu Asp
115 120 125

Asn Glu Arg Ser Asn Gly Thr Ile Ile His Ile Lys Glu Lys His Leu
130 135 140

35

Cys His Thr Xaa Xaa Xaa Gln Ser Ser Pro Lys Leu Phe Trp Ala Leu
145 150 155 160

Tyr Val Val Ala Gly Val Leu Phe Cys Tyr Gly Leu Leu Val Thr Val
165 170 175

40

Ala Leu Cys Val Ile Trp Thr Asn Ser Arg Arg Asn Arg Leu Leu Gln
180 185 190

~~Val Thr Tyr Met Asn Met Thr Pro Arg Arg Pro Gly Leu Thr Arg Xaa~~
~~195 200 205~~

45

Lys Pro Tyr Gln Pro Tyr Ala Pro Ala Arg Asp Phe Ala Ala Tyr Arg
210 215 220

Pro
225

50

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 225 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

5 Met Thr Leu Arg Leu Leu Phe Leu Ala Leu Ser Phe Phe Xaa Ser Val
1 5 10 15

Gln Val Thr Glu Asn Lys Ile Leu Val Lys Gln Ser Pro Leu Leu Val
20 25 30

10 Tyr Asp Asn Asn Glu Val Xaa Ser Leu Ser Cys Arg Tyr Ser Tyr Asn
35 40 45

Leu Leu Ala Lys Glu Phe Arg Ala Ser Leu Tyr Lys Gly Val Asn Ser
50 55 60

15 Asp Val Xaa Glu Val Cys Val Gly Asn Gly Asn Phe Thr Tyr Gln Pro
65 70 75 80

Gln Phe Arg Pro Asn Val Gly Phe Asn Cys Asp Gly Asn Phe Asp Asn
85 90 95

20 Glu Thr Val Thr Phe Arg Leu Trp Asn Leu Asp Val Asn His Thr Asp
100 105 110

Ile Tyr Phe Cys Lys Ile Glu Val Met Tyr Pro Pro Pro Tyr Leu Asp
115 120 125

25 Asn Glu Lys Ser Asn Gly Thr Ile Ile His Ile Lys Glu Lys His Leu
130 135 140

Cys His Ala Xaa Xaa Xaa Gln Thr Ser Pro Lys Leu Phe Trp Pro Leu
145 150 155 160

30 Val Val Val Ala Gly Val Leu Leu Cys Tyr Gly Leu Leu Tyr Thr Val
165 170 175

Thr Leu Cys Ile Ile Trp Thr Asn Ser Arg Arg Asn Arg Leu Leu Gln
180 185 190

Ser Asp Tyr Met Asn Met Thr Pro Arg Arg Leu Gly Pro Thr Arg Xaa
195 200 205

35 Lys His Tyr Gln Pro Tyr Ala Pro Ala Arg Asp Phe Ala Ala Tyr Arg
210 215 220

Pro
225

(2) INFORMATION FOR SEQ ID NO:5:

~~(i) SEQUENCE CHARACTERISTICS:~~

- (A) LENGTH: 223 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

50 Met Leu Arg Leu Leu Leu Ala Leu Asn Leu Phe Pro Ser Ile Gln Val
1 5 10 15

Thr Gly Asn Lys Ile Leu Val Lys Gln Ser Pro Met Leu Val Ala Tyr
20 25 30

55 Asp Asn Ala Tyr Xaa Asn Leu Ser Cys Lys Tyr Ser Tyr Asn Leu Phe

35 40 45
 Ser Arg Glu Phe Arg Ala Ser Leu His Lys Gly Leu Asp Ser Ala Val
 50 55 60
 5
 Xaa Glu Val Cys Val Val Tyr Gly Asn Tyr Ser Gln Gln Leu Gln Val
 65 70 75 80
 Tyr Ser Lys Thr Gly Phe Asn Cys Asp Gly Lys Leu Gly Asn Glu Ser
 85 90 95
 10
 Val Thr Phe Tyr Leu Gln Asn Leu Tyr Val Asn Gln Thr Asp Ile Tyr
 100 105 110
 Phe Cys Lys Ile Glu Val Met Tyr Pro Pro Pro Tyr Leu Asp Asn Glu
 115 120 125
 15
 Lys Ser Asn Gly Thr Ile Ile His Val Lys Gly Lys His Leu Cys Pro
 130 135 140
 Ser Pro Leu Phe Pro Gly Pro Ser Lys Pro Phe Trp Val Leu Val Val
 145 150 155 160
 20
 Val Gly Gly Val Leu Ala Cys Tyr Ser Leu Leu Tyr Thr Val Ala Phe
 165 170 175
 Ile Ile Phe Trp Val Arg Ser Lys Arg Ser Arg Leu Leu His Ser Asp
 180 185 190
 25
 Tyr Met Asn Met Thr Pro Arg Arg Pro Gly Pro Thr Arg Xaa Lys His
 195 200 205
 Tyr Gln Pro Tyr Ala Pro Pro Arg Asp Phe Ala Ala Tyr Arg Ser
 210 215 220

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 226 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

40 Met. Leu Gly Ile Leu Val Val Leu Cys Leu Ile Pro Ala Ala Asp Val
 1 5 10 15
 Thr Glu Asn Lys Ile Leu Val Ala Gln Arg Pro Leu Leu Ile Val Ala
 20 25 30
 45 Asn Arg Thr Ala Xaa Thr Leu Val Cys Asn Tyr Thr Tyr Asn Gly Thr
 35 40 45
 Gly Lys Glu Phe Arg Ala Ser Leu His Lys Gly Thr Asp Ser Ala Val
 50 55 60
 50 Xaa Glu Val Cys Phe Ile Ser Trp Asn Met Thr Xaa Lys Ile Asn Ser
 65 70 75 80
 Asn Ser Asn Lys Glu Phe Asn Cys Arg Gly Ile His Asp Lys Asp Lys
 85 90 95
 55 Val Ile Phe Asn Leu Trp Asn Met Ser Ala Ser Gln Thr Asp Ile Tyr
 100 105 110

Phe Cys Lys Ile Glu Ala Met Tyr Pro Pro Pro Tyr Val Tyr Asn Glu
 115 120 125
 5 Lys Ser Asn Gly Thr Val Ile His Tyr Arg Glu Thr Pro Ile Xaa Xaa
 130 135 140
 Gln Thr Gln Glu Pro Glu Ser Ala Thr Ser Tyr Trp Val Met Tyr Ala
 145 150 155 160
 10 Val Thr Gly Leu Leu Gly Phe Tyr Ser Met Leu Ile Thr Ala Val Phe
 165 170 175
 Ile Ile Tyr Arg Gln Lys Ser Lys Arg Asn Arg Tyr Arg Gln Ser Asp
 180 185 190
 15 Tyr Met Asn Met Thr Pro Arg His Pro Pro His Gln Lys Asn Lys Gly
 195 200 205
 Tyr Pro Ser Tyr Ala Pro Thr Arg Asp Tyr Thr Ala Tyr Arg Ser Trp
 210 215 220
 20 Gln Pro
 225

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: protein
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:
 Met Tyr Pro Pro Ala Tyr
 1 5

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 6 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: protein
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Tyr Pro Pro Pro Ala
 1 5

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: cDNA
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

AATACGACTC ACTATAGG

18

- 5 (2) INFORMATION FOR SEQ ID NO:10:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - 10 (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: cDNA
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

15 CACCACACTG TATTAACC

18

- (2) INFORMATION FOR SEQ ID NO:11:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 6 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: protein
 - 25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:
Met Tyr Pro Pro Pro Tyr
1 5

- 30 (2) INFORMATION FOR SEQ ID NO:12:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 base pairs
 - (B) TYPE: nucleic acid
 - 35 (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: cDNA
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

40 TGCAAGGTGG AGCTCATGTT CCCACCGCCA TAC

33

-
- (2) INFORMATION FOR SEQ ID NO:13:
- 45 (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - 50 (ii) MOLECULE TYPE: cDNA
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GCGCTCGACT CTAGAAGCAT CCTCGTG

27

55

Claims

1. A CTLA4 mutant molecule reactive with the CD80 antigen, wherein in an extracellular domain of CTLA4 the first tyrosine in the amino acid motif MYPPPY is replaced by an amino acid other than tyrosine.
2. The CTLA4 mutant molecule of claim 1, wherein the amino acid replacing the first tyrosine is alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, or valine.
3. The CTLA4 mutant molecule of claim 1, wherein the amino acid replacing the first tyrosine is phenylalanine.
4. The CTLA4 mutant molecule of claim 1, wherein the amino acid replacing the first tyrosine is tryptophan.
5. A functional, soluble CTLA4 mutant molecule of claim 1 or 2.
6. The CTLA4 mutant molecule of claim 6 which binds the CD80 antigen but not the CD86 antigen.
7. The CTLA4 mutant molecule of claim 6 joined to a non-CTLA4 molecule.
8. The CTLA4 mutant molecule of claim 7, wherein the non-CTLA4 protein is at least a portion of an immunoglobulin molecule.
9. The CTLA4 mutant molecule of claim 1, wherein the molecule is a CTLA4Ig mutant fusion molecule having a first amino acid sequence containing amino acid residues corresponding to the extracellular domain of CTLA4 mutant molecule and a second amino acid sequence containing amino acid residues corresponding to the hinge, CH2 and CH3 regions of immunoglobulin C γ l.
10. A purified protein of claim 1 or 9.
11. A nucleic acid molecule encoding the protein of claim 10.
12. A DNA molecule of claim 11.
13. A cDNA molecule of claim 12.
14. A plasmid encoding the cDNA molecule of claim 13.
15. A host-vector system comprising the plasmid of claim 14 transfected into a compatible eucaryotic host cell.
16. The host-vector system of claim 15, wherein the compatible eucaryotic host cells are Chinese Hamster Ovary cells.
17. A method for producing a CTLA4 mutant molecule comprising growing the host vector system of claim 15 so as to produce the CTLA4 mutant molecule in the host and recovering the CTLA4 mutant molecule so produced.
18. A method for regulating an immune response by blocking a CD80 interaction with lymphocytes comprising contacting a CD80 positive cell with the CTLA4 mutant molecule of claim 1 to interfere with reaction of endogenous CD80 antigen with CTLA4.
19. The method of claim 18, wherein the lymphocytes are CD80 positive lymphocytes.
20. The method of claim 19, wherein the immune response is a B cell response resulting in the inhibition of antibody production, a T cell response resulting in inhibition of cell mediated immunity is an inhibition of lymphocyte proliferation.
21. A method for treating immune system diseases mediated by T cell interactions with CD80 positive cells comprising administering to a subject the CTLA4 mutant molecule of claim 1, to regulate T cell interactions with said CD80 positive cells.

22. A CTLA4 mutant molecule reactive with the CD80 antigen, wherein the amino acid residue at the position corresponding to tyrosine - 99 in wild type CTLA4 is not tyrosine or alanine.

5

10

15

20

25

30

35

40

45

50

55

FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7

Wild type (w.t.) and Mutant Binding to CD80 CHO and CD86 CHO
as determined by FACS




FIGURE 8

MYPPPY (wild type) and mutant Binding to B7-1 CHO and B7-2 CHO
as determined by FACS





FIGURE 9

MYPPPY (w.t.) and MFPPPY (1 ug/mL) Binding to LCL 816 Cells
in the Presence of anti-CD80 or anti-CD86
as Determined by FACS



(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 757 099 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
26.05.1999 Bulletin 1999/21

(51) Int Cl.⁶: C12N 15/12, C07K 14/705,
C12N 15/85, C12N 5/10,
A61K 38/17

(43) Date of publication A2:
05.02.1997 Bulletin 1997/06

(21) Application number: 96305298.0

(22) Date of filing: 19.07.1996

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE

• Ledbetter, Jeffrey A.
Seattle, WA 98117 (US)
• Peach, Robert
Edmonds, WA 98026 (US)

(30) Priority: 21.07.1995 US 505058

(71) Applicant: Bristol-Myers Squibb Company
New York, N.Y. 10154 (US)

(74) Representative: Jones, Alan John et al
CARPMAELS & RANSFORD
43 Bloomsbury Square
London, WC1A 2RA (GB)

(72) Inventors:
• Linsley, Peter S.
Seattle, WA 98119 (US)


(54) CTLA4 mutant molecules and uses thereof

(57) The invention provides CTLA4 mutant molecules as ligands for the B7 antigen. Methods are provided for expressing CTLA4 mutant molecules as soluble,

functional molecules, for preparing CTLA4 mutant fusion proteins, and for using these soluble molecules to regulate T cell interactions and immune responses mediated by such interactions.

FIGURE 7

Wild type (wt) and mutant Binding to CD80 CHO and CD86 CHO
as determined by FACS



EP 0 757 099 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 96 30 5298

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	LINSLEY P S ET AL: "Human B7-1 (CD80) and B7-2 (CD86) bind with similar avidities but distinct kinetics to CD28 and CTLA-4 receptors." IMMUNITY, (1994 DEC) 1 (9) 793-801. JOURNAL CODE: CCF. ISSN: 1074-7613., XP002098031 United States * the whole document *	1,2,5,6, 9,11-17	C12N15/12 C07K14/705 C12N15/85 C12N5/10 A61K38/17
X	PEACH R J ET AL: "COMPLEMENTARITY DETERMINING REGION 1 (CDR1)- AND CDR3-ANALOGOUS REGIONS IN CTLA-4 AND CD28 DETERMINE THE BINDING TO B7-1" JOURNAL OF EXPERIMENTAL MEDICINE, vol. 180, no. 6, 1 December 1994, pages 2049-2058, XP000199779 * the whole document *	1,2,5,9, 11-17	
A	WO 93 00431 A (SQUIBB BRISTOL MYERS CO) 7 January 1993 * the whole document *		TECHNICAL FIELDS SEARCHED (Int.Cl.6)
X,P	EP 0 682 039 A (SQUIBB BRISTOL MYERS CO) 15 November 1995 * claim 19; example 7 *	1,2,5,9, 11-17	C07K C12N
X,P	MORTON P A ET AL: "Differential effects of CTLA-4 substitutions on the binding of human CD80 (B7-1) and CD86 (B7-2 binding of human)." JOURNAL OF IMMUNOLOGY, (1996 FEB 1) 156 (3) 1047-54. JOURNAL CODE: IFB. ISSN: 0022-1767., XP002098032 United States * the whole document *	1,2,5,6, 9,11-17	

The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 25 March 1999	Examiner: Bilang, J
CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document			

EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 96 30 5298

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

25-03-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9300431 A	07-01-1993	AT 170562 T	15-09-1998
		AU 661854 B	10-08-1995
		AU 2240092 A	25-01-1993
		CA 2110518 A	07-01-1993
		DE 69226871 D	08-10-1998
		EP 0606217 A	20-07-1994
		ES 2123001 T	01-01-1999
		FI 935795 A	22-12-1993
		JP 6508989 T	13-10-1994
		MX 9203605 A	01-11-1993
		NO 934801 A	21-02-1994
		NZ 243286 A	26-03-1996
		NZ 264712 A	26-03-1996
		PT 100637 A	31-05-1994
		US 5770197 A	23-06-1998
		US 5434131 A	18-07-1995
		US 5844095 A	01-12-1998
		US 5851795 A	22-12-1998
		US 5885796 A	23-03-1999
		US 5885579 A	23-03-1999
		ZA 9204782 A	27-12-1993
EP 0682039 A	15-11-1995	AU 701310 B	28-01-1999
		AU 1645895 A	26-10-1995
		CA 2146895 A	16-10-1995
		FI 951801 A	16-10-1995
		JP 8047391 A	20-02-1996
		NO 951436 A	16-10-1995
		US 5844095 A	01-12-1998
		US 5851795 A	22-12-1998
		US 5885796 A	23-03-1999
		US 5773253 A	30-06-1998
		US 5885579 A	23-03-1999

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82